

# International Journal of Clinical Obstetrics and Gynaecology

ISSN (P): 2522-6614  
ISSN (E): 2522-6622  
© Gynaecology Journal  
[www.gynaecologyjournal.com](http://www.gynaecologyjournal.com)  
2022; 6(2): 86-87  
Received: 07-01-2022  
Accepted: 09-02-2022

## Rahul Sakpal

Assistant Professor, Department of Pathology, BKL Walawalkar Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Supriya Bapat

Postgraduate Student, Department of Obstetrics and Gynaecology, BKL Walawalkar Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Samrudhini Kapale

Senior Resident, Department of Obstetrics and Gynaecology, BKL Walawalkar Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Dr. Abhijit Ambike

Professor, Department of Obstetrics and Gynaecology, BKL Walawalkar Rural Medical College and Hospital, Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Unmesh Santpur

Professor & Head, Department of Obstetrics and Gynaecology, BKL Walawalkar Rural Medical College and Hospital, Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Deepali Ambike

Professor & Head, Department of Paediatrics, PCMC, S Post Graduate Institute, YCM hospital, Pimpri, Pune, Maharashtra, India

## Corresponding Author:

### Dr. Abhijit Ambike

Professor, Department of Obstetrics and Gynaecology, BKL Walawalkar Rural Medical College and Hospital, Dervan, At Post Sawarda, Taluka Chiplun, Ratnagiri, Maharashtra, India

## Struma ovarii: A case report

Rahul Sakpal, Supriya Bapat, Samrudhini Kapale, Dr. Abhijit Ambike, Unmesh Santpur and Deepali Ambike

DOI: <https://doi.org/10.33545/gynae.2022.v6.i2b.1167>

### Abstract

**Background:** Struma ovarii, an ovarian tumor, is diagnosed when >50% of the teratoma is thyroid tissue. Struma ovarii's incidence is <2% of mature teratoma.

**Case:** We present a rare case report of struma ovarii of a 47-year-old female who presented with incomplete evacuation of bladder and bowel since 2 years. She also complained of dragging pain in abdomen, intermittent radiating to back since last 2/3 years. Her carcino embryonic antigen (CEA) level was 2.56 ng/mL and histologically proved to be struma ovarii.

**Results:** Single cystic 7 × 4 cm left ovarian mass was diagnosed as struma ovarii.

**Conclusion:** Struma ovarii though a rare entity should always be kept in mind in a pelvic mass lesion causing discomfort. It is not usually a malignant condition and histopathological examination should be done to rule out struma ovarii.

**Keywords:** G2P2, tubectomised, abdomen

### Introduction

Struma ovarii is a rare and usually unilateral specialized or monodermal teratoma <sup>[1]</sup>. It is predominantly composed of mature thyroid tissue <sup>[2]</sup>. It accounts for approximately 5 percent of all ovarian teratomas <sup>[3-5]</sup>. Struma ovarii is most common between the ages of 40 and 60 years <sup>[5]</sup>. Women with struma ovarii usually present with pain and/or a pelvic mass and less frequently with ascites. Clinical and biochemical features of hyperthyroidism are uncommon in women with struma ovarii, occurring in less than 5 to 8 percent of cases <sup>[4, 6-9]</sup>.

### Case Report


A 47-year-old female, farmer by occupation and no addictions referred by general practitioner who was G2P2, tubectomised with complaints of incomplete evacuation of bladder and bowel since last 2/3 years and increased frequency and urgency for urination. She had generalised pain in abdomen, dragging in nature, radiating to back, left more than right side intermittently since 2-3 months and bloating sensation in abdomen with pressure on rectum. Pain relieved temporarily on rest and medication from local doctor and aggravated by movement. Patient c/o dyspareunia intermittently. No excessive p/v bleeding, or white p/v discharge.

She was married since 28 years with two alive children and normal menstrual cycles. No significant medical, surgical or family history. On per abdominal examination it was soft, nontender, and mass of 12-14 weeks palpable. firm in consistency, external ballotment absent.

Both ultrasonography and CT-scan report commented as left ovarian neoplasm. Hysterectomy with salpingo-oophorectomy was done and later it was diagnosed struma ovarii (left) by histopathological examination.



**Fig 1:** Gross Specimen: Single cystic 7 × 4 cm left ovarian mass



**Fig 2:** Histopathology: showed thyroid tissue comprised of varying sizes follicles and without any other germ cell origin tissue

### Discussion

Struma ovarii is a monodermal teratoma <sup>[1]</sup>. Struma ovarii accounts for approximately 5 percent of all ovarian teratomas <sup>[3-5]</sup>. The thyroid gland typically is not enlarged, abdominal pain was present in 20 to 42 percent and a palpable lower abdominal mass in 23 to 58 percent <sup>[5, 9]</sup>. Struma ovarii cannot be diagnosed on clinical, biochemical or imaging test. Diagnosis is confirmed by Histopathological examination. Microscopically, it shows mature thyroid tissue <sup>[2]</sup>. Removal of the lesion, commonly by excision is the preferred treatment.

### Conclusion

Although it is a pelvic mass lesion and causes discomfort, it is not usually a malignant condition and histopathological examination should be done to rule out struma ovarii.

**Consent:** A written informed consent was obtained from her for publication of this case report and its accompanying images

**Contributors:** AA, US: manuscript writing, patient management; RS, SK, SB: manuscript editing, patient management.

### References

1. Robins & Cotran Pathologic Basis of Disease. Editor Kumar, Abbas, Fausto. 9th edition, page 1030; 2015, Publisher Elsevier India Ltd.
2. Dunzendorfer T, de Las Morenas A, Kalir T, Levin RM.

- Struma ovarii and hyperthyroidism. *Thyroid*. 1999;9:499.
3. Kondi-Pafiti A, Mayrigiannaki P, Grigoriadid Ch, *et al*. Monodermal teratomas (Struma ovarii). Clinicopathological characteristics of 11 cases and literature review. *Eur J Gynaecol Oncol*. 2011;32:657.
4. Yassa L, Sadow P, Marqusee E. Malignant struma ovarii. *Nat Clin Pract Endocrinol Metab*. 2008;4:469.
5. Yoo SC, Chang KH, Lyu MO, *et al*. Clinical characteristics of struma ovarii. *J Gynecol Oncol*. 2008;19:135.
6. Kraemer B, Grischke EM, Staebler A, *et al*. Laparoscopic excision of malignant struma ovarii and 1 year follow-up without further treatment. *Fertil Steril*. 2011;95:2124.e9.
7. Ayhan A, Yanik F, Tuncer R, *et al*. Struma ovarii. *Int J Gynaecol Obstet*. 1993;42:143.
8. Young RH. New and unusual aspects of ovarian germ cell tumors. *Am J Surg Pathol*. 1993;17:1210.
9. De Simone CP, Lele SM, Modesitt SC. Malignant struma ovarii: a case report and analysis of cases reported in the literature with focus on survival and 1131 therapy. *Gynecol Oncol*. 2003;89:543