

# International Journal of Clinical Obstetrics and Gynaecology

ISSN (P): 2522-6614  
ISSN (E): 2522-6622  
Indexing: Embase  
Impact Factor (RJIF): 6.71  
© Gynaecology Journal  
[www.gynaecologyjournal.com](http://www.gynaecologyjournal.com)  
2025; 9(6): 1205-1209  
Received: 02-08-2025  
Accepted: 05-09-2025

**Dr. Monika Rahar**  
Senior Resident, Department of  
Obstetrics and Gynaecology, PDU  
Medical College, Churu, Rajasthan,  
India

**Dr. Surendra Singh**  
Assistant Professor, Department of  
General Medicine, SMS Medical  
College and Hospital, Jaipur,  
Rajasthan, India

**Dr. Jyoti Ola**  
Senior Resident, Department of  
Obstetrics and Gynaecology, JLN  
Medical College, Ajmer, Rajasthan,  
India



## Severe maternal anemia presenting in late gestation: Its influence on delivery in a tertiary care centre in South-East Rajasthan

Monika Rahar, Surendra Singh and Jyoti Ola

DOI: <https://www.doi.org/10.33545/gynae.2025.v9.i6g.1784>

### Abstract

**Background:** Anaemia is the most common nutritional deficiency in the world. Globally, anaemia affects 1.62 billion people which constitute to 24.8% of the total population and the group with the greatest number of individuals affected being pregnant women (41.8%). The condition is prominent in Southeast Asian countries where about half of all global maternal deaths are due to anemia and India contributes about 80% of the maternal death due to anaemia in South Asia. There is marginally decrease in prevalence of anemia in pregnant women in India from 58% in NFHS-3 (National Family Health Survey-2005-06) to 50% in NFHS-4 survey (2015-16).

**Material and Methods:** This is a prospective study conducted for six months from February 2021 to July 2021 in the Department of Obstetrics and Gynaecology at a tertiary care hospital, Jhalawar. Data was collected from bedside medical records. A total of 190 women were studied who fulfilled the inclusion and exclusion criteria.

**Results:** A total of 4911 deliveries occurred during the study period, of which 190 (3.87%) women had severe anemia. Most women (59.5%) were younger than 25 years. A majority were of low socioeconomic status, illiterate, unbooked, and multiparous. LSCS was the commonest mode of delivery (58.4%). Maternal complications were reported in 98 women (51.6%). Preterm labour was most common (35.8%), preeclampsia occurred in 14.2%, antepartum hemorrhage in 10.5%, postpartum hemorrhage in 8.9%, abruptio placentae in 8.9%, IUGR in 7.4%, sepsis in 2.6%, and placenta previa in 6.3%. Neonatal complications occurred in 87 cases (45.8%). NICU admission was required in 41.6%, prematurity occurred in 21.6%, respiratory distress in 12.1%, meconium-stained liquor in 9.5%, and intrauterine fetal demise in 4.2%.

**Conclusion:** Severe anemia in pregnancy significantly increases the risk of complications during labour and delivery. Improving antenatal care coverage, promoting iron-folic acid supplementation, raising nutritional awareness, addressing parasitic infections, and strengthening rural healthcare infrastructure can reduce maternal and neonatal morbidity. Preventive strategies, timely interventions, and public health initiatives targeting women of reproductive age are essential to reduce the impact of severe anemia on fetomaternal health.

**Keywords:** Gestational thrombocytopenia, pregnancy, asymptomatic thrombocytopenia

### Introduction

Anaemia is the most common nutritional deficiency in the world. Globally, anaemia affects 1.62 billion people which constitute to 24.8% of the total population and the group with the greatest number of individuals affected being pregnant women (41.8%). The condition is prominent in Southeast Asian countries where about half of all global maternal deaths are due to anemia and India contributes about 80% of the maternal death due to anaemia in South Asia. There is marginally decrease in prevalence of anemia in pregnant women in India from 58% in NFHS-3 (National Family Health Survey-2005-06) to 50% in NFHS-4 survey (2015-16).

Anaemia in pregnancy is one of the most common medical problem encountered by obstetricians, especially in developing countries. World Health Organization has reported, 35% to 75% (56% on average) of pregnant women in developing countries and 18% of women from industrialized countries are anemic [2]. Many of these women were already anemic at the time of conception. The prevalence of anemia is high in central Asia and has been reported as 62-85% in India [3, 4, 5, 6].

The centre of disease control and prevention (1990) defined anaemia as less than 11 gm/dl in first and third trimester and less than 10.5 gm/dl in second trimester.7

**Corresponding Author:**  
**Dr. Monika Rahar**  
Senior Resident, Department of  
Obstetrics and Gynaecology, PDU  
Medical College, Churu, Rajasthan,  
India

**Table 1:** WHO defines anaemia in pregnancy as hemoglobin concentration of less than 11 gm% (7.45 mmol/L) and hematocrit less than 33% (.8)

Category	Anaemia severity	Hb level in gm/dl
1	Mild	9-10.9
2	Moderate	7-8.9
3	Severe	<7

**Table 2:** Anaemia is further classified into mild, moderate and severe by ICMR depending upon the levels of hemoglobin (.9)

Category	Anaemia severity	Hb level in gm/dl
1	Mild	10-10.9
2	Moderate	7-9.9
3	Severe	4-6.9
4	Very severe	<4

Severe anemia is defined as hemoglobin levels less than 7 gm/dL [4]. These women have decreased oxygen carrying capacity of blood. Severe anemia in women who are already in labour is a critical situation for the obstetrician with the risk of adverse feto-maternal outcomes. It is responsible for 20-40% of direct and indirect maternal deaths because of increased susceptibility to cardiac failure, sepsis and association with preeclampsia, antepartum haemorrhage, postpartum haemorrhage and thrombo-embolism [3, 10 11]. Risk of preterm delivery, low birth weight, prematurity, intrauterine growth retardation, intrauterine death and birth asphyxia is increased causing increased perinatal morbidity and mortality in neonate [10, 12, 13].

Anemia has multifactorial etiology .Nutritional anemia is more common i.e. inhibitors of iron absorption, dietary deficiency of iron, folic acid and vitamin B12. During pregnancy, fetal and placental growth and larger amount of circulatory blood leads to the increased demand for nutrients, especially iron and folic acid. Other factors are pregnancy iron deficiency, teenage pregnancy, lack of appropriate spacing between pregnancies, parasitic infestation (e.g. malaria, hookworm infestation), poor environmental and personal hygiene

### Aim and objectives

To study women with gestational age beyond 28 weeks presenting in the emergency OPD with severe anemia in labour and to find out its feto-maternal outcome and negative impact on fetus and maternal health status.

### Material and Methods

This is prospective study conducted for six months from

February 2021 to July 2021 in the Department of Obstetrics and Gynaecology at tertiary care hospital Jhalawar. Data was collected from bed side medical records. 190 women were studied, they were fulfilling inclusion and exclusion criteria.

### Inclusion criteria

- Patients in labour with gestational age>28 weeks with severe anemia (Hb< 7 g/dl)
- Singleton pregnancy

### Exclusion criteria

- Patients with low haemoglobin levels whose pregnancy <28 weeks excluded from the study.
- Women with chronic illness like-kidney disease, liver disease.
- Patients with hemoglobinopathy- like sickle cell anemia, thalassemia.
- Multiple pregnancy
- Women with bone marrow disorder.
- Women with bleeding disorder
- Patients whose pregnancy >28 weeks gestational age but haemoglobin levels >7 g/dl.

Medical records of patients fulfilling the inclusion were studied for maternal age, parity, booking status, gestational age, preeclampsia, low birth weight, NICU admission, fetal demise, postpartum hemorrhage, maternal intensive care, maternal mortality.

### Results

Number of deliveries during the study period was 4911. Out of 4911 deliveries 190 (3.87%) were severely anaemic. Maximum women were < 25 years of age group (59.5%), Hindu (90%), illiterate (56.3%), House wife (63.7%), unbooked (67.4%), low socioeconomic status (49.5%) and multipara (60%) . Mostly delivered by LSCS (58.4%).

In our study, out of the 190 participants, maximum 68 had preterm labour (35.8%) followed by 27 (14.2%) had preeclampsia, APH was found in 20 (10.5%), PPH in 17 (8.9%), IUGR in 14 (7.4%), abruptio placentae in 17 (8.9%), placenta previa in 12 (6.3%) and sepsis in 5 (2.6%) participants.

In our study, out of the 190 participants, most common fetal complication was prematurity in 41(21.6%), NICU admission was required in 79 (41.6%) and respiratory distress was found in 23(12.1%), MSL in 18 (9.5%) and IUD was in 8 (4.2%) participants.


**Table 3:** Sociodemographic and Obstetric Characteristics of the Study Population (n = 190)

Characteristics	Frequency	No. of women
<b>Age group</b>		
≤ 25 years	113	59.5
25-30 years	53	27.9
31-35 years	14	7.4
>35 years	10	5.3
<b>Religion</b>		
Hindu	171	90.0
Muslim	19	10.0
<b>Education</b>		
Illiterate	107	56.3
Primary school	40	21.1
Secondary school	28	14.7
Graduate and above	15	7.9
<b>Occupation</b>		
Farmer	21	11.1
House wife	121	63.7


Labor	30	15.8
Student	18	9.5
<b>Socioeconomic status</b>		
Lower	94	49.5
Middle	88	46.3
Upper	8	4.2
<b>Booking status</b>		
Unbooked	128	67.4
Booked	62	32.6
<b>Parity</b>		
Primi	76	40.0
Multi	114	60.0
<b>Mode of delivery</b>		
LSCS	111	58.4
Vaginal	79	41.6
<b>Gestational age</b>		
28-32 weeks	15	7.9
33-36 weeks	74	38.9
>37	101	53.2

**Table 4:** Maternal Complications among Women with Severe Anemia (n = 190)

<b>Maternal complications</b>	<b>Frequency</b>	<b>Percent</b>
Preterm labour	68	35.8
PE	27	14.2
APH	20	10.5
PPH	17	8.9
IUGR	14	7.4
Sepsis	5	2.6
Abruptio placentae	17	8.9
Placenta previa	12	6.3

**Fig 1:** Distribution of participants according to maternal complications

<b>Fetal complications</b>	<b>Frequency</b>	<b>Percent</b>
NICU admission	79	41.6
Prematurity	41	21.6
Respiratory distress	23	12.1
MSL	18	9.5
IUD	8	4.2



**Fig 2:** Distribution of Participants According to Fetal Complications

## Discussion

In India it is not uncommon to see patients with severe anaemia late in pregnancy with no prior antenatal visits especially in low socioeconomic settings and the same is evident from our study. Anemia in pregnancy is the major health issue in rural part of India due to various reasons like poverty, illiteracy, lack of awareness about the need for antenatal care and presence of superadded infections. Anemia in pregnancy is the risk factor for preterm labour, intrauterine growth restriction, cardiac failure, puerperal sepsis, sub-involution and failure of lactation. Severe anemia in pregnancy has a significant impact on maternal and perinatal outcomes. Severity of anemia is an independent risk factor for preeclampsia, low birth weight.

In our study, mean age of study participants was  $25.3 \pm 5.7$  years and out of the 190 participants, maximum 113 (59.5%) were < 25 years age followed by 53 (27.9%) in age of 25-30 years, 14 (7.4%) in age of 31 - 35 years and 10 (5.3%) in age of more than 35 years.

A similar study was done by Singh S *et al* (18), in this study a total 210 deliveries with severe anaemia were included and maximum participants were in age of 20-24 years (42.8%).

In our study, out of the 190 participants, 171 (90%) were Hindu participants and 19 (10%) were Muslim participants. this distribution could be explained on the bases of population ratio. Study by Singh R *et al* also found similar kind of results and most of the participants were belongs to Hindu religion.

In this study, out of the 190 participants, maximum 107 (56.3%) were illiterate followed by 40 (21.1%) were educated up to primary school, maximum 121 (63.7%) were housewife followed by 30 (15.8%) were engaged in labor work and maximum 94 (49.5%) belongs to lower socioeconomic status followed by 88 (46.3%) belongs to middle class. This distribution could be explained because our centre is a tertiary care centre and most of the study participants are rural, referred and unbooked. The cause of high prevalence of anaemia could be explained on basis of poor nutrition because of lack of awareness regarding ANC care.

Study by Singh S *et al* revealed that maximum women belonged to low socioeconomic status (87.14%), living in rural area (72.8%).

In our study, out of the 190 participants, 128 (67.4%) were unbooked pregnancy, 114 (60%) were multipara, 111 (58.4%) delivered by LSCS and 79 (41.6%) by vaginal delivery, and mean gestational age at the time of delivery was  $36.6 \pm 2.8$  weeks

and maximum 101 (53.2%) had gestational age >37 weeks followed by 74 (38.9%) had 33-36 weeks at the time of delivery and 15 (7.9%) had 28-32 weeks.

A similar study was done by Singh S *et al*. in this study multipara were 60.4% and maximum 57.6% had gestational age >37 at the time of delivery.

In this study, mean Hb of study participants was  $6.4 \pm 0.56$  gm%. In our study, out of the 190 participants, 98 (51.6%) had any kind of maternal morbidity and maximum 68 (35.8%) had preterm labour followed by 27 (14.2%) had preeclampsia, APH was found in 20 (10.5%), PPH in 17 (8.9%), IUGR in 14 (7.4%), abruptio placentae in 17 (8.9%) and placenta previa found in 12 (6.3%) participants.

Singh S *et al* revealed that commonest complication of severe anaemia observed in the present study was preterm labour showing highest incidence of 89 cases (42.8%) followed by preeclampsia (16.1%), intrauterine growth restriction (9.0%), intrauterine death (8.5%), sepsis (3.8%), abruptio placentae in 3.3%, placenta previa (2.3%), cardiac failure in 1.4%. Maternal mortality was seen only in one case (0.47%). Cause of maternal mortality was decompensated cardiac failure due to severe anaemia superimposed with severe preeclampsia. Anaemia prevalence was also significantly high in pregnant women from low socioeconomic status (87.6%) compared to those from middle socioeconomic status (12.4%).

In this study, out of the 190 participants, neonatal mortality was found to be in 10 (5.3%) and neonatal morbidity was found to be in 87 (45.8%) and among them NICU admission was required in 79 (41.6%) and prematurity was found in 41(21.6%), respiratory distress was found in 23 (12.1%), MSL in 18 (9.5%) and IUD was in 8 (4.2%) participants.

Ruchika Singh *et al* also did a similar study and found that Mean birth weight was more in without anemia group (p value 0.002). Birth asphyxia and NICU admissions were more in severe anemia group neonates (p value 0.012, 0.017 respectively).

## Conflict of Interest

Not available

## Financial Support

Not available

## Conclusion

Anemia in pregnancy is a major health problem in developing

countries, contributing significantly to fetomaternal morbidity and mortality. Proper Antenatal care should be made accessible and affordable to both urban and rural areas. Proper antenatal counselling regarding diet, birth spacing, regular antenatal checkups and regular intake of iron supplements to be done in all antenatal clinics. By keeping this in view, it is recommended that good antenatal care should be made available, accessible and affordable to all pregnant women through partnership between all tiers of government and non-governmental organizations. New and innovative strategies are needed, particularly those that improve the overall health and nutrition status of adolescent girls before they enter their reproductive years.

## References

1. Milman N. Postpartum anemia I: definition, prevalence, causes and consequences. *Ann Haematol.* 2011;90:1247-1253.
2. World Health Organization. The global prevalence of anaemia in 2011. Geneva: World Health Organization; 2015.
3. Suryanarayana R, Chandrappa M, Santhuram AN, Prathima S, Sheela SR. Prospective study on prevalence of anemia of pregnant women and its outcome: a community based study. *J Fam Med Prim Care.* 2017;6(4):739-743.
4. Agarwal KN, Agrawal DK, Sharma A, Sharma K, Prasad K, Kalita MC, et al. Prevalence of anemia in pregnant and lactating women in India. *Food Nutr Bull.* 2006;27:311-315.
5. World Health Organization. The prevalence of anemia in pregnancy. WHO Technical Report Series. Geneva: World Health Organization; 1992-1993.
6. Prashant D, Jaideep KC, Girija A, Mallapur MD. Prevalence of anemia among pregnant women attending antenatal clinics in rural field practice area of Jawaharlal Nehru Medical College, Belagavi, Karnataka, India. *Int J Community Med Public Health.* 2017;4(2):537-541.
7. Centers for Disease Control and Prevention. Criteria for anemia in childbearing age group. *MMWR.* 1989;38:400-404.
8. World Health Organization. Vitamin and Mineral Nutrition Information System. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. Geneva: World Health Organization; 2011.
9. Indian Council of Medical Research. Evaluation of nutritional anaemia prophylaxis programme: task force study. New Delhi: Indian Council of Medical Research; 1989.
10. Maka SS, Tondare SB, Tondare MB. Study of impact of anemia on pregnancy. *Int J Reprod Contracept Obstet Gynecol.* 2017;6(11):4847-4850.
11. Singh S, Kaur K. Maternal outcome in cases of severe anaemia in labour. *Int J Reprod Contracept Obstet Gynecol.* 2018;7(3):1201-1204.
12. Sangeetha VB, Pushpalatha S. Severe maternal anemia and neonatal outcome. *Sch J Appl Med Sci.* 2014;2(1C):303-309.
13. Marhatta R. Study of anemia in pregnancy and its outcome in Nepal Medical College Teaching Hospital, Kathmandu, Nepal. *Nepal Med Coll J.* 2007;9:270-274.
14. Bansal P, et al. Women with severe anemia in labor: fetomaternal outcomes. *Int J Health Sci Res.* 2022;12(1):2249-9571.
15. Sabina S, Iftequar S, Zaheer Z, Khan M, Khan S. An overview of anemia in pregnancy. *JIPBS.* 2015;2:144-151.
16. Karaoglu L, Pehlivan E, Egri M, Deprem C, Gunes G, Genc MF, et al. The prevalence of nutritional anemia in pregnancy in an east Anatolian province, Turkey. *BMC Public Health.* 2010;10:329-335.
17. Singh S, Kaur K. Maternal outcome in cases of severe anemia in labour. *Int J Reprod Contracept Obstet Gynecol.* 2018;7(3):1201-1204.
18. Singh R, Gupta M, Saini V. Women with severe anemia in labour: adverse clinical outcomes. *Int J Innov Res Med Sci.* 2019;4(1):2455-8737.

### How to Cite This Article

Rahar M, Singh S, Ola J. Severe maternal anemia presenting in late gestation: Its influence on delivery in a tertiary care centre in South-East Rajasthan. *International Journal of Clinical Obstetrics and Gynaecology* 2024; 9(6): 1205-1209.

### Creative Commons (CC) License

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International (CC BY-NC-SA 4.0) License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.