

# International Journal of Clinical Obstetrics and Gynaecology

ISSN (P): 2522-6614  
ISSN (E): 2522-6622  
© Gynaecology Journal  
[www.gynaecologyjournal.com](http://www.gynaecologyjournal.com)  
2020; 4(4): 65-67  
Received: 20-11-2019  
Accepted: 23-12-2019

**Dr. Nabnita Patnaik**  
Associate Professor, Department of  
Obstetrics & Gynaecology, SVS  
Medical College and Hospital,  
Mahbubnagar, Telangana, India

**Dr. C Venkatrami Reddy**  
Professor, Department of  
Obstetrics & Gynaecology  
SVS Medical College and Hospital,  
Mahbubnagar, Telangana, India

**Dr. RL Tejaswi Tirumala**  
Assistant Professor, Department of  
Obstetrics & Gynaecology, SVS  
Medical College and Hospital,  
Mahbubnagar, Telangana, India

**Dr. Chapay Soren**  
Professor and Head, Department  
of Paediatrics, SVS Medical College  
and Hospital, Mahbubnagar,  
Telangana, India

**Dr. Mandepudi Geethika**  
Assistant Professor, Department of  
Radiodiagnosis, SVS Medical  
College and Hospital,  
Mahbubnagar, Telangana, India

**Corresponding Author:**  
**Dr. Nabnita Patnaik**  
Associate Professor, Department of  
Obstetrics & Gynaecology, SVS  
Medical College and Hospital,  
Mahbubnagar, Telangana, India

## Outcome of pregnancies with fibroids and its associated complications: A prospective study

**Dr. Nabnita Patnaik, Dr. C Venkatrami Reddy, Dr. RL Tejaswi Tirumala,  
Dr. Chapay Soren and Dr. Mandepudi Geethika**

**DOI:** <https://doi.org/10.33545/gynae.2020.v4.i4b.621>

### Abstract

**Aim:** To assess the outcome of pregnancies with fibroids and associated complications.

**Materials and Methods:** The present prospective observational clinical study was conducted in the Department of Obstetrics and Gynecology, SVS Medical College and Hospital, MBNR. From October-2017 to September-2019, 110 cases of were found to be eligible for inclusion in the study.

**Results:** Mean age of the study subjects was 27.24 years. 33.6% patients were primigravida and 66.4% were multigravida. 70.0% patients reported single fibroid and 30.0% reported multiple fibroids. 26 (23.6%) had spontaneous abortion

**Conclusion:** Fibroids were observed to be responsible for high incidence of complications throughout pregnancy and during postpartum period.

**Keywords:** Maternal outcome, Fibroid, Spontaneous abortion, Gravidity

### Introduction

Myomas are the most frequently recorded benign smooth muscle tumor of the uterus, affecting 20%–60% of women of reproductive age [1]. At present, although there are a lot of research about the prevention and treatment of uterine fibroids the etiopathogenesis of uterine fibroids is still unclear. The incidence of fibroids in pregnancy reported ranges from 0.1 to 10.7% of all pregnancies and increases as the female chooses to postpone pregnancy later on [2]. It was found that 10%–40% of prepartum complications which happened in pregnancy with fibroid have been associated with the presence of it [3]. Also, they are related to a lot of ante-, intra-, and postpartum complications [4].

There are conflicting data on the relationship between obstetric outcomes and uterine fibroids, and the mechanism by which fibroids influence obstetric outcomes is unclear. Some studies have shown a relationship between uterine fibroids and pregnancy complications, such as preterm birth, premature rupture of membranes (PROM), fetal malpresentation, placental abruption and intrauterine fetal demise [5-7].

In addition, uterine fibroids have been linked to labor dystocia, puerperal infection, operative vaginal delivery, cesarean delivery and postpartum hemorrhage (PPH) [8]. In contrast; other studies have reported no increased risks for these adverse obstetric outcomes with uterine fibroids [9].

Hence the present study was conducted with the aim to assess the outcome of pregnancies with fibroids and any associated complications.

### Materials and Methods

The present prospective observational clinical study was conducted in the Department of Obstetrics and Gynecology, SVS Medical College and Hospital, MBNR, From October-2017 to September-2019.

### Inclusion Criteria

1. Those who have provided the informed consent
2. Singleton pregnancy
3. Gestational age of 24-42 weeks at the time of delivery
4. Patients with fibroid of  $\geq 2$  cm were included in the study

### Exclusion Criteria

1. Women with pathological conditions (chronic hypertension, gestational diabetes or pre-existing diabetes mellitus, uterine anomalies or fetal malformations).
2. Multiple- pregnancy
3. Patients who have not signed the informed consent

### Ethical approval and Informed consent

The study protocol was reviewed by the Ethical Committee of the Hospital and granted ethical clearance. After explaining the purpose and details of the study, a written informed consent was obtained.

### Data collection

Maternal demographic characteristics, medical and obstetrical history and pregnancy outcomes were collected from Medical Record Viewer database of Hospital along with manual retrieval from medical charts and labor records using standardized data collection forms. Of the 588 women who delivered during the study period, 110 patients met the inclusion criteria.

### Statistical analysis

The recorded data was compiled and entered in a spreadsheet computer program (Microsoft Excel 2010) and then exported to data editor page of SPSS version 19 (SPSS Inc., Chicago, Illinois, USA). Descriptive statistics included computation of percentages and means.

### Results

**Table 1:** Demographic and clinical profile of the study population

Age	27.24±5.26
BMI	26.19±4.39
Gravida	
Primigravida	37 (33.6%)
Multi-gravida	73 (66.4%)
Type of Conception	
Spontaneous	58 (52.8%)
Assisted	52 (47.2%)
Number of Fibroids	
Single	77 (70.0%)
Multiple	33 (30.0%)

**Table 2:** Antenatal and postnatal maternal outcome

Outcome	N (%)
Threatened Miscarriage	16 (14.5%)
Pre-term Labor	13 (11.8%)
Placenta Previa	3 (2.7%)
Postpartum hemorrhage	2 (1.79%)
Spontaneous abortion	26 (23.6%)
Premature delivery	16 (14.5%)
Cesarean sections	40 (36.4%)

**Table 3:** Fetal Outcome

Outcome	
Fetal Weight (Kgs)	2.71±0.76
NICU stay (Days)	2.31±1.39
Congenital anomaly	1 (0.91%)

### Discussion

Mean maternal age in our study was found to be 27.24 years, which is comparable to other studies, showing occurrence of leiomyomas in second and third decades of life<sup>[10, 11]</sup>.

In the present investigation 33.4% patients were primigravida

and 66.4% were multigravida. We found that fibroids were less frequent in first pregnancy compared to multigravida. This is in consistent with earlier studies conducted by Noor *et al.*<sup>[11]</sup> and Sarwar *et al.*<sup>[10]</sup>

Regarding obstetric complications, in our study 26 (23.6%) had spontaneous abortion. The proposed mechanism is compressed endometrial vascular supply, affects the fetus adversely resulting in abortion<sup>[10]</sup>.

In the present investigation 70.0% patients reported single fibroid and 30.0% reported multiple fibroids. Lam *et al.*<sup>[12]</sup> reported a higher rate of preterm delivery among patients with multiple fibroids compared with those with a single fibroid. Likewise, Ciavattini *et al.*<sup>[7]</sup> monitored raised preterm delivery, cesarean delivery, and breech presentation rates among individuals with multiple fibroids compared with single fibroids or no fibroids. However, Lai *et al.*<sup>[9]</sup> recorded no relationship between preterm delivery and fibroid number.

In our study, cesarean section was reported among 36.4% of the subjects. Similarly in various studies, rate of cesarean section ranges 34%-73%. Klatsky *et al.* 2008 recorded that women with fibroids were at a 3.7-fold increased risk of cesarean delivery<sup>[13]</sup>. Our study had limitations of being just observational one not having a comparing group, the sample size was small, and some popular concepts could have resulted in a high cesarean delivery rate.

### Conclusion

Most of the fibroids are asymptomatic but may adversely affect the path of pregnancy and labor dependent on their location and size. The present study revealed that fibroids were found to be responsible for high incidence of complications throughout antepartum, intrapartum, and postpartum period. So, they have to be carefully screened in the antenatal period through regular follow-up.

### References

1. Laughlin SK, Schroeder JC, Baird DD. New directions in the epidemiology of uterine fibroids. *Seminars in Reproductive Medicine.* 2010; 28(3):204-17.
2. Somigliana E, Vercellini P, Daguati R. Fibroids and female reproduction: A critical analysis of the evidence. *Human Reproduction Update.* 2007; 3(5):465-76.
3. Eze CU, Odumeru EA, Ochie K, Nwadike UI, Agwuna KK. Sonographic assessment of pregnancy co-existing with uterine leiomyoma in Owerri, Nigeria. *African Health Sciences.* 2013; 13(2):453-60.
4. Stout MJ, Odibo AO, Graseck AS, Macones GA, Crane JP, Cahill AG. Leiomyomas at routine second-trimester ultrasound examination and adverse obstetric outcomes. *Obstet Gynecol.* 2010; 116(5):1056-63.
5. Qidwai GI, Caughey AB, Jacoby AF. Obstetric outcomes in women with sonographically identified uterine leiomyomata. *Obstet Gynecol.* 2006; 107(2 Pt 1):376-82.
6. Ezzedine D, Norwitz ER. Are Women With Uterine Fibroids at Increased Risk for Adverse Pregnancy Outcome? *Clin Obstet Gynecol.* 2016; 59(1):119-27.
7. Ciavattini A, Clemente N, Delli Carpini G, Di Giuseppe J, Giannubilo SR, Tranquilli AL. Number and size of uterine fibroids and obstetric outcomes. *J Matern Fetal Neonatal Med.* 2015; 28(4):484-88.
8. Vergani P, Locatelli A, Ghidini A, Andreani M, Sala F, Pezzullo JC. Large uterine leiomyomata and risk of cesarean delivery. *Obstet Gynecol.* 2007; 109(2 Pt 1):410-14.

9. Lai J, Caughey AB, Qidwai GI, Jacoby AF. Neonatal outcomes in women with sonographically identified uterine leiomyomata. *J Matern Fetal Neonatal Med.* 2012; 25(6):710-713.
10. Sarwar I, Habib S, Bibi A, Malik N, Parveen Z. Clinical audit of foetomaternal outcome in pregnancies with fibroid uterus. *J Ayub Med Coll Abbottabad.* 2012; 24(1):79-82.
11. Noor S, Fawwad A, Sultana R, *et al.* Pregnancy with fibroids and its and its obstetric complication. *J Ayub Med Coll Abbottabad.* 2009; 21(4):37-40.
12. Lam SJ, Best S, Kumar S. The impact of fibroid characteristics on pregnancy outcome. *Am J Obstet Gynecol.* 2014; 211(4):395.e1-395.
13. Klatsky PC, Tran ND, Caughey AB, Fujimoto VY. Fibroids and reproductive outcomes: a systematic literature review from conception to delivery. *Am J Obstet Gynecol.* 2008; 198(4):357-66.