

International Journal of Clinical Obstetrics and Gynaecology

ISSN (P): 2522-6614
ISSN (E): 2522-6622
© Gynaecology Journal
www.gynaecologyjournal.com
2021; 5(2): 121-124
Received: 05-01-2021
Accepted: 08-02-2021

Rehana Rashid

Senior Resident, Department of
Obstetrics and Gynaecology, GMC,
Srinagar, Jammu and Kashmir,
India

Ufaque Muzaffar

Lecturer, Department of Obstetrics
and Gynaecology, GMC, Srinagar,
Jammu and Kashmir, India

Javid Ahmad Sofi

Senior Resident, Department of
Clinical Haematology, SKIMS,
Srinagar, Jammu and Kashmir,
India

Zohra Younus

Senior Resident, Department of
Obstetrics and Gynaecology, Jamia
Hamdard, Delhi, India

Corresponding Author:

Rehana Rashid

Senior Resident, Department of
Obstetrics and Gynaecology, GMC,
Srinagar, Jammu and Kashmir,
India

Fetomaternal outcome in cases of severe anemia in labour

Rehana Rashid, Ufaque Muzaffar, Javid Ahmad Sofi and Zohra Younus

DOI: <https://doi.org/10.33545/gynae.2021.v5.i2c.874>

Abstract

Introduction: Anaemia is a major public health problem throughout the world especially in developing countries like India and it is the most common nutritional deficiency disorder in the world [1]. Anemia in pregnancy is defined as Hemoglobin concentrations <11 g/dL in the 1st and last trimester, and a Hemoglobin concentration of <10.5 g/dL in the 2nd trimester. The prevalence of anaemia indicates the nutritional status of a community and it often goes untreated especially in pregnant women where it can be most dangerous [4].

Materials and Methods: The present study was a prospective observational study conducted at Post Graduate Department of Obstetrics and Gynaecology LallaDed hospital Srinagar. The study was conducted over a time period of 12 months from December 2019 to December 2020. The study included 70 patients with anemia and 70 patients without anemia.

Results: The mean age of patients in anemia group was 30.63±4.97 years and the mean age in control group was 30.60±5.17 years. However the *p* value was statistically insignificant (0.97). Regarding antenatal registration, maximum patients 30.7% in anemia group were unbooked and 19.3% were booked. While as in group B, 27% were booked and 23% patients were unbooked. 31.4% patients from anemia group and 27% patients from group B were multigravida and 18.6% patients from anemia group and 23% patients from group B were primigravida. The difference wasn't statistically significant (*p* value 0.39). The inter pregnancy interval in Anemia group was <2 years in 31.4% patients and >2 years in 18.6% patients while as in group B, maximum patients (40%) had inter pregnancy interval of >2 years and only 10% had <2 years. The difference between two groups regarding inter pregnancy interval was statistically significant (*p* value<0.05).

Conclusion: Anemia in pregnancy is a major health problem in developing countries, contributing significantly to fetomaternal morbidity and mortality. Proper Antenatal care should be made accessible and affordable to both urban and rural areas. Proper antenatal counselling regarding diet, birthspacing, regular ante natal checkups and regular intake of iron supplements to be done in all antenatal clinics.

Keywords: Anemia, pregnancy, PPH, antenatal

Introduction

Anaemia is a major public health problem throughout the world especially in developing countries like India and it is the most common nutritional deficiency disorder in the world [1]. World Health Organization (WHO) has estimated the prevalence of anaemia among pregnant population to be about 38% [2]. Anemia in pregnancy is defined as Hemoglobin concentrations <11 g/dL in the 1st and last trimester, and a Hemoglobin concentrations of <10.5 g/dL in the 2nd trimester [3]. The prevalence of anaemia indicates the nutritional status of a community and it often goes untreated especially in pregnant women where it can be most dangerous [4].

Iron deficiency anaemia is an important public health problem for pregnant women, living in developing countries, affecting 2/3rd of pregnant women and contributes to maternal morbidity and mortality and to low birth weight [5, 6]. Iron deficiency can be due to inadequate oral intake or poor bioavailability of iron in foodstuffs. It can also occur due to excessive loss of iron from the body. While the diet may contain adequate amounts of iron poor bioavailability of dietary iron is considered as the major reason for the widespread prevalence of iron deficiency anaemia [7-9]. Folate deficiency anaemia has been seen in 25-50% of pregnant women in some areas of India attending the hospital clinics [7]. Studies have shown that folate deficiency anaemia in pregnancy has become a global problem and mainly affects the economically unprivileged population [10-16]. Anaemia has hazardous influence on maternal and fetal outcome and increases risk of postpartum haemorrhage (PPH), infection, sepsis and risk for pre-term birth,

low birth weight and small for gestational age babies thereby contributing to maternal and perinatal morbidity and sometimes mortality [17]. In India, anaemia is the second most common cause of maternal death, accounting for 20% of total maternal deaths [15]. Considering the effects of IDA on maternal and fetal mortalities, physical function and child growth and development, it is regarded as one of the main health indicators [4]. In fact the high incidence of post-partum haemorrhage (PPH) in India as compared to rest of world has been attributed to the high prevalence of anaemia in pregnant women [2].

Aims and Objectives

- To study the role of age and other demographic factors on anemia.
- To evaluate the fetomaternal outcome due to anemia in pregnancy.

Materials and Methods

The present study was a prospective observational study conducted at Post Graduate Department of Obstetrics and Gynaecology LallaDed hospital Srinagar. The study was conducted over a time period of 12 months from December 2019 to December 2020. The study included 70 patients with anemia and 70 patients without anemia.

Inclusion criteria

Singleton pregnancy, Pregnant women beyond 36 weeks of gestation, HB<7.

Exclusion criteria

Multiple pregnancy, Patients with anemia due to acute blood loss, Women with chronic medical illness.

Procedure

Seventy pregnant women with hb<7, reporting to OPD or admitted to the hospital in labour or not in labour after 37 weeks of pregnancy were taken as study subjects. Seventy pregnant women without anemia admitted to the hospital were taken as control group. All the haematological and biochemical investigations were carried out on admission (Complete hemogram, Peripheral blood smear, Routine urine examination, Iron profile, Ultrasonography for fetal well being). Patients were decided for caesarean sections or were induced for labour as per the obstetric indications. Patients were monitored for incidence of intrapartum, postpartum and perinatal complications.

Statistical analysis

Statistical testing was conducted with the statistical package for the social science system version (SPSS). Continuous variables were expressed as Mean \pm SD and categorical variables were summarized as frequencies. The statistical significance of the difference between two groups was based on p-value. P-value of <0.05 was considered to be statistically significant.

Observations and Results

In this study 70 patients with anemia have been compared to 70 patients without anemia to evaluate the role of age, sociodemographic factors, parity on anemia and to study the influence of anemia on fetomaternal unit. The mean age of patients in anemia group was 30.63 \pm 4.97 years and the mean age in control group was 30.60 \pm 5.17 years. However the p-value was statistically insignificant (0.97). The mean gestational age in group A was 37.40 \pm 1.02 and 37.26 \pm 0.943 in group B. The p-value was 0.393. In anemia group, 30% patients belonged to

Rural area and 20% patients belonged to Urban area where as in control group 22% patients belonged to rural areas and 27% belonged to Urban areas. The p-value was statistically insignificant (0.23). Regarding antenatal registration, maximum patients 30.7% in anemia group were unbooked and 19.3% were booked. While as in group B, 27% were booked and 23% patients were unbooked. However the difference was statistically insignificant (p value 0.90). Regarding parity, 31.4% patients from anemia group and 27% patients from group B were multigravida and 18.6% patients from anemia group and 23% patients from group B were primigravida. The difference wasn't statistically significant (p value 0.39). The inter pregnancy interval in Anemia group was <2 years in 31.4% patients and >2 years in 18.6% patients while as in group B, maximum patients (40%) had inter pregnancy interval of >2 years and only 10% had <2 years. The difference between two groups regarding inter pregnancy interval was statistically significant (p value<0.05). The mode of delivery was FTVD in 31% patients from anemia group and 36% patients from non-anemia group, LSCS in 19% patients from anemia group and 14% from non-anemia group (p-value=1).




Fig 1: Anemia in rural and urban patients




Fig 2: Anemia in multi vs primi

Regarding maternal complications, from anemia group out of 70 patients, 3(2.1%) had Abruptio, 5(3.6%) had Preterm labour, 7(5%) had postpartum haemorrhage and only 1(0.7%) had sepsis. In non-anemic group, 2(1.4%) had Abruptio, 2(1.4%) had Preterm labour, 4(2.9%) had post-partum haemorrhage and none had sepsis. However the p value was statistically insignificant (0.16). The maximum babies in anemia group had APGAR score of 7(23.6%) followed by 8(14.3%). In non-anemia group maximum babies had APGAR score of 8(23.6%) followed by 7(18.6%). The p value was 0.087. In anemia group patients out of 70 babies, 3(2.1%) were IUGR, 1(0.7%) had IUFD, 5(3.6%) had meconium aspiration and 4(2.9%) had respiratory distress syndrome. In non anemiagroup, only 2(1.4%) had meconium aspiration and 4(2.9%) had respiratory distress syndrome. The p value was statistically insignificant (0.240).




Fig 3: Effect of inter-pregnancy interval




Fig 4: APGAR score

Discussion

Anemia in pregnancy is more prevalent and severe in developing countries [17]. Maternal anemia has been found to be associated with higher maternal mortality and morbidity and adverse perinatal outcome [18, 19]. Older women have more burden of anemia than younger ones, this might be due to combined effects of repeated pregnancies, inadequate dietary intake and social

factors. In our study, the majority of pregnant women were in age group of 30-34 years. Regular antenatal checkups and access to the health facilities are cornerstone for the safe motherhood. In our study, 30.7% patients from anemia group were unbooked and 30% patients were from rural areas. Anemia in pregnancy is more common in women of high parity and birth spacing of <2 years [20]. In our study 31.4% patients with anemia were multi gravida and 31.4% patients had inter pregnancy interval of <2 years. There was no significant difference between the modes of deliveries in both groups. Maternal effects of anaemia vary, depending upon the degree of anaemia. It can directly lead to cardiac failure, haemorrhage and infection. Severe anaemia may impair myometrial contractility resulting from impaired oxygen carrying capacity leading to enzyme and cellular dysfunction which leads to postpartum haemorrhage (PPH) [21]. In our study, presence of anemia was associated with occurrence of PPH, Abruptio, preterm labour and sepsis. However the p value was statistically insignificant (0.169). Anaemia in pregnancy is associated with adverse foetal outcome in the form of preterm birth, low birth weight, intrauterine growth restriction, stillbirths, intrauterine foetal death, and birth asphyxia. In the present study the incidence of IUGR was significantly associated with anemia. The incidence of meconium aspiration and respiratory distress syndrome was also high in anemic patients.

Conclusion

Anemia in pregnancy is a major health problem in developing countries, contributing significantly to fetomaternal morbidity and mortality. Proper Antenatal care should be made accessible and affordable to both urban and rural areas. Proper antenatal counselling regarding diet, birth spacing, regular ante natal checkups and regular intake of iron supplements to be done in all antenatal clinics.

References

1. Kaur K. Anaemia 'a silent killer' among women in India: Present scenario. *European Journal of Zoological Res* 2014;3(1):32-6.
2. The global prevalence of anaemia in 2011. World Health Organization; 2015. appswho.int/iris/bitstream/10665/177094/1/9789241564960_eng.pdf.
3. WHO. Centers for Disease Control and Prevention Atlanta. Worldwide Prevalence of anemia 1993–2005. [www.who.int. http://whqlibdoc.who.int/publications/2008/9789241596657_eng.pdf](http://www.who.int/http://whqlibdoc.who.int/publications/2008/9789241596657_eng.pdf). Published 2008.
4. Panghal R, Boora P. Prevalence of anaemia among pregnant women of low-income group of Hisar district of Haryana. *J Dairying, Foods Home Sci* 2010;29(2):112-5.
5. Pasricha SR, Caruana SR, Phuc TQ, Casey GJ, Jolley D, Kingsland S *et al*. Anemia, iron deficiency, meat consumption, and hookworm infection in women of reproductive age in northwest Vietnam. *Am J Trop Medicine and Hygiene* 2008;78(3):375-81.
6. Baig-Ansari N, Badruddin SH, Karmaliani R, Harris H, Jehan I, Pasha O *et al*. Anemia prevalence and risk factors in pregnant women in an urban area of Pakistan. *Food Nutri Bulletin* 2008;29(2):132-9.
7. WHO/UNICEF/UNU. Iron Deficiency Anemia: Assessment, Prevention and Control. Geneva, Switzerland: World Health Organization 2001.
8. Narasinga Rao BS, Vijayasathya C, Prabhavathi T. Iron absorption from habitual diets of Indians studied by the extrinsic tag technique. *Indian J Med Res* 1983;77:648-57.

9. Nadiger HA, Krishnamachari KAVR, Nadamuni Naidu A, NarasingaRao BS, Srikantia SG. The use of common salt (sodium chloride) fortified with iron to control anemia: results of a preliminary study. *Br J Nutr* 1980;43:45–51.
10. World Health Organization (WHO). The prevalence of Anaemia in women: a tabulation of available information. Geneva, Switzerland: WHO 1992.
11. Luwang NC, Gupta VM, Khanna S. Anaemia in pregnancy in rural community of Varanasi. *Ind J Prev Soc Med* 1980;11:83-3.
12. Agrawal V, Tejwanin S. Prevalence of iron deficiency anaemia in Indian antenatal women especially in rural areas. *Ind Med Gaz* 1999, 300-3.
13. National Family Health Survey for India conducted by. Mumbai, India: International Institute for Population Science 2006.
14. De Benoist B, McLean E, Egli I, Cogswell M. WHO global database on anaemia. Geneva: WHO; 2008. Worldwide prevalence of anaemia 1993-2005.
15. Ezzati M, Lopez AD, Dogers A, Vander HS, Murray C. Selected major risk factors and global and regional burden of disease. *Lancet* 2002;360:1347-60.
16. Irshad G, Jafri SA, Kousar S, Ali I. Significance of serum ferritin in diagnosis of iron deficiency anaemia in pregnant females of Pakistan. *Professional Med J* 2011;18(3):475-78.
17. Diejomaeh FME, Abdulaziz A, Adekile AD. Anemia in pregnancy. *Int J Obstet Gynecol* 1999;65(3):299-301.
18. Bhatt R. Maternal mortality in India FOGSI-WHO study. *J Obstet Gynecol Ind* 1997;47:207-14.
19. Reddy V, Rao NP, Sastry JG, Kashinath K. Nutritional trends in India. Hyderabad National Institute of Nutrition 1993.
20. Konar H. DC Dutta's Textbook of Obstetrics. JP Medical Ltd 2014.
21. Tusimin M, Yazit A, Zainulddin NS, Vaiappuri VSSKN, Md Noor S. The Impact of Severity of Antenatal Anaemia on Maternal and Perinatal Outcome in Hospital Serdang, Central Malaysia. *J Preg Child Health* 2016;3:291.